



Microbes and Infectious Diseases

Journal homepage: <https://mid.journals.ekb.eg/>

Original article

Knowledge mapping of Nigeria's scientific contribution to antimicrobial resistance research: A visualized investigation using VOS viewer and cite space

Shuaibu Suleiman Adeiza^{*1}, Mohammed Gazali Shuaibu², Abdulmalik Bello Shuaibu³

1- Department of Pharmaceutical Microbiology, Faculty of Pharmaceutical sciences, Ahmadu Bello University, Zaria, Kaduna, Nigeria.

2- Department of Microbiology, Faculty of Chemical and Life sciences, Usmanu Danfodiyo University, Sokoto, Nigeria.

3- Department of Veterinary Microbiology, Faculty of Veterinary medicine, Usmanu Danfodiyo University, Sokoto, Nigeria..

ARTICLE INFO

Article history:

Received 2 October 2022

Received in revised form 10 November 2022

Accepted 11 November 2022

Keywords:

Antimicrobial resistance
Bibliometric
Antimicrobial stewardship
Drug discovery
Nigeria

ABSTRACT

Aim: To identify the contribution and impact of Nigerian authors, institutions, and journal sources, assess the knowledge base, detect the trends regarding antimicrobial resistance research through bibliometric analysis. **Methods:** Data from a query of antimicrobial resistance (AMR) articles in the dimensions database (PubMed included) were subjected to bibliometric analysis (1972–2022). Bibliometric indicators, topic networks were established, and topic trends were analysed by frequency, relevance scores, and trends over time. **Results and conclusion:** A total of 0.2% of the papers on AMR published worldwide were written by authors or institutions from Nigeria. In 2021, publications grew by 13.6%. The vast majority of publications (57.18%) were in the field of medical and health sciences, with 2428 papers. The top journal for AMR papers with at least one Nigerian institution affiliation was African Journal of Clinical and Experimental Microbiology, which published 1.8% of these papers. Antimicrobial stewardship, clinical and laboratory practices on AMR, public health implications, traditional and molecular methods, and phytomedicine and drug discovery were prominent areas of focus for Nigerian researchers. This study provided a comprehensive overview of the AMR research output, highlighting the need of increased research capacity to address the burden of AMR in Nigeria. Because Nigeria is overly reliant on conventional methods of discovery and development, innovation is essential for the future of antibiotics.

Introduction

Microbes can become resistant to antimicrobials through mutation or adaptation, which renders treatments ineffective. This phenomenon is known as antimicrobial resistance (AMR) [1]. The World Health Organization (WHO) estimates that AMR kills over 700,000 people across the globe yearly, an estimate that may well soar to 10 million by the year 2050 [2]. According to a needs assessment on antimicrobial

use and resistance in Nigeria published in 2017 by the federal ministries of health, agriculture, and the environment, certain antimicrobial agents have increased resistance rates of 70% to 100% in certain states [3]. In a country where infectious diseases are a major cause of morbidity and mortality, Nigeria considers AMR a priority on the national public health agenda[3]. The Nigerian response to antimicrobial resistance led by the Nigeria Centre

for Disease Control (NCDC) began in 2016 following a situational analysis that investigated common antimicrobial-resistant pathogens recovered from hospitals, animal sources, agricultural and environmental sources[4]. Overall, the National Action Plan (NAP) strives to address five key pillars in accordance with the WHO Global Action Plan on AMR [5]. The fundamentals entail boosting knowledge and awareness of AMR among health personnel and the general public; developing a one health surveillance system; strengthening prevention efforts, control, and biosecurity; promoting rational use of antimicrobials and access to quality medications; and conducting research into antimicrobial alternatives, new diagnostics, and drug discovery[4]. The problem of AMR in Nigeria is multisectoral. The ease with which antibiotics are available and can be purchased without the need for a prescription thrives in an environment where the antibiotic market is poorly regulated. The problem of improper disposal of expired antibiotics and left-over medicines by the general public, which mostly end up in conventional waste bins and dumps due to lack of awareness on the part of the users and weak enforcement on the part from regulatory authorities[6]. The problem of indiscriminate use impacts human health and animal welfare alike [7]. In the human health sector, many incidences of fever are addressed empirically with antibiotics[8]. Although a few antimicrobials are prescribed by a doctor, patients and their families frequently press healthcare professionals for antimicrobials and, in most cases, self-medicate with antimicrobials even for mild viral infections[9]. Even with the Nigerian veterinary council's warnings against excessive use of antimicrobials in the animal health sector, several farmers habitually add antimicrobials into animal feed or water to promote growth and, in the case of poultry farming, to boost egg size to ensure better profit. Several farmers in the country use antimicrobials in post-harvest storage to preserve the yield[10]. As a result, human and animal waste, particles from crop sprays, discharge from health facilities, pharmaceutical companies, and farmlands carrying antimicrobial-resistant pathogenic microbes and antimicrobial residues find their way into the ecosystem[11]. It thus results in successions of negative impact on food production, bodies of water, as well as fish farms[11]. Despite increased awareness of antimicrobial resistance, the general situation is deteriorating, and the scientific community and Institutions have endured in

research and publications to fashion newer antimicrobial peptides (AMPs), antibiotic combinations, and monitoring systems to effectively control AMR[12]. The amount of reliance on data increases with the growth of scientific publications and information technologies in healthcare. These big data raise various issues to be resolved by innovative big data analytics, like Knowledge mapping via bibliometrics.

Knowledge mapping is an assessment tool that blends concepts and techniques from mathematics, visuals, and other disciplines with bibliometric methods[13]. Cite space, and visualization of similarities (VOS) are the most popular bibliometric analysis tool for knowledge mapping via bibliometrics[14]. Knowledge mapping has been applied in many research fields[13, 15, 16]. To date, a knowledge mapping analysis of AMR research contributions and inclinations in Nigeria has remained hidden. This study aims to identify the contribution and impact of Nigerian authors, institutions, and journal sources, assess the knowledge base, detect the trends regarding antimicrobial resistance research through bibliometric analysis. Understanding Nigeria's contribution to AMR research requires quantifying and mapping AMR-related publications. This study examines the research conducted by institutions in Nigeria using the Dimensions database as a foundation. This descriptive study focuses on bibliometric elements like international collaboration, subject area classification, active authors and institutions, journals and highly cited papers. In this study, we generated visualized knowledge maps of AMR research in Nigeria and analysed scientific hotspots, emerging/future trends, prolific authors and key research institutions of momentous research activities in AMR, which we hope will provide clarity about the nation's priority for AMR research; and serve as reference for future collaborative networks.

Methods

Data acquisition and query criteria

Bibliographic data on antimicrobial resistance publications authored by a Nigerian researcher were retrieved from the Dimensions™ database on August 20th 2022. The database is one of the most comprehensive for bibliometric and patentometric analysis. It contains millions of publications cited over 1.6 billion times linked with patents, clinical trials, policy documents, datasets and grants [15, 17]. This database is completely inclusive of

PubMed, making it highly reliable. We used the search criteria: 'antibiotics OR antifungal OR antibacterial OR antimicrobial resistance AND Nigeria' to spotlight research articles published between 1972 and 2022. The search was not limited by language, and it included all accessible data sets. Both title and abstracts were searched to ensure subject-specific outcome for the benefit of accuracy. Publications indexed in the database that featured a minimum of one author connected to a Nigerian institution, including a university or organization, qualified for inclusion.

Data processing and parsing

The dplyr and ggplot2 packages of R- studio v.4.1.2 (2022-02-16) was used to examine, parse and visualize bibliographic data. Imported raw files were cured of false positives and duplicates. Top 100 documents were arbitrarily selected per year and checked for presence of false positive results. A key term was ascribed to noun-phrases with similar meanings (e.g., "Antimicrobial sensitivity test", and "AST" were merged to "antimicrobial susceptibility test"). Multiple or similar keyword occurrences in a publication were consolidated as a single count.

Network mapping

Network mapping was rendered with visualization of similarities (VOS) viewer software (Leiden University, Netherlands)[18]. This method adopts diverse clusters and color spectrum; the bubble size symbolizes the total frequency, the colours indicate the cluster type, and the lines of conjunct (straight or curved) illustrates connectedness [19]. Indices like, scientific production, mean citations, authors productivity, affiliation and countries, preferred journals and keywords, were used to assess the present and likely future trends of AMR research in Nigeria [20, 21]. The terms used in keyword clustering analysis were extracted from title and abstract fields of included bibliographic documents. Copyright statements and structured abstract labels were ignored in the analysis. Full counting method with minimum occurrence threshold of 20 ($T \geq 20$) was used. The top 60% of calculated relevance scores were used in the analysis.

Citation burst and centrality index

In order to determine the burstness and centrality index of the bibliographic components, cite space (Version 6.1.R3 Basic) was used[22]. The following limits were set: time span (1972 to 2022), years per slice (one), pruning (pathfinder and Pruning Sliced Networks), selection criteria (g-index), link strength (cosine) and scope (within slices) and other default settings.

Furthermore, the top cited articles were also extracted and sorted by field citation ratio. The field citation ratio (FCR) and relative citation ratio (RCR) are citation-based measures of an articles' scholarly impact[23]. FCR was determined by dividing the

number of citations received by an article by average number of citations received by articles published within the same year and fields of research (FOR) class. While the RCR is measured as the number of citations received by a paper adjusted to the citation count garnered by National Institute of Health (NIH) funded publications in the same research field and year[23].

Results

Nigeria's annual contribution in AMR Research

Nigerian authors and institutions have played a key role in production of a total of 4,246 papers on antimicrobial resistance. As noted in **figure (1)**, AMR-related research in Nigeria have shown an annual upward tendency that peaked in 2021(13.6%). Based on our research, there are 2,176,583 publications on AMR globally and Nigeria have contributed at least 0.2% of that number. **Table S1** shows the distribution of scientific output by field of research; medical and health sciences (n = 2428, 57.18%), biological sciences (n = 806, 18.98%), agricultural and veterinary sciences (n = 409, 9.63%).

Influential contributor analyses

1. Authors and journals contributing to AMR research

A total of 14391 Nigerian authors were involved in AMR research. OKEKE IN published the most papers indexed in Dimension database (n = 31), followed by LAMIKANRA A (n = 28), and OLAYINKA BO (n = 27) (**Figure 2a**). By analysing the distribution of journals accepting Nigeria's contribution in this field, we could accurately identify the main part of the academic research, and papers published in such journals. **Figure 2b** shows the top journals that published Nigeria's contributions in AMR research. The number of publications in these journals was about 21.83% of the total number of contributions in this field. The number of articles published in african journal of clinical and experimental microbiology was the largest, with 75 articles, accounting for 1.8% of the total. Followed by PLoS One and african journal of medicine and medical sciences, with 50 articles each (1.2%). In terms of mean citation, American Journal of Tropical Medicine and Hygiene had the highest (33.3%).

The top cited author by centrality is BAUER AW (1985) with centrality of 0.13 followed by Okeke IN (2000), with centrality of 0.07 (**Figure 3a**).

2. Contributions and cooperation network among institutions/organizations

The top ranked institutions/organizations in AMR research collaboration by centrality are University of Ibadan, with centrality of 0.33. The second one was Ahmadu Bello University, with centrality of 0.14. The third was University of Lagos, with centrality of 0.13. The 4th was International Institute of Tropical Agriculture, with centrality of 0.13. The 5th was Obafemi Awolowo University, with centrality of 0.10 (**Figure 3b**).

A co-authorship network map constructed based on authors (n = 1000) who had published at least five papers on AMR is shown in **Figure (4)**. In the figure, authors that share the same colour space represent the same cluster. There were active collaborations in AMR research in Nigeria, especially among authors in the same cluster, such as OKEKE IN AND LAMIKANRA A (green), SOWUNMI A and ODUOLA AMJ (coffee brown), ILIYASU G and HABIB AG (lime green), OLAYINKA BO AND ONAOLAPO JA (yellow) etc.

Authors with citation bursts are defined as those that are cited frequently over a while. Through detecting bursts, **Table 1** lists the top authors of publications on AMR with the strongest citation bursts. COKER A., led the list with the maximum burst duration of 15 years and strength of 4.54, followed by OGUNSOLA F with a 14-year citation burst with 4.45 burst strength. NGWAI Y had a minimum burst duration of 1 year and the highest strength of 8.7. Recently published authors of AMR research with the strongest burst were NGWAI Y (8.7), OLAYINKA B (8.53) and OKEKE IN (7.27). This shows that their work may have formed a hot and leading topic. The table also show a list of the top cited authors of publications in AMR with the strongest citation bursts. AKO-NAI AK on the top of the list with the maximum burst duration of 18 years and strength of 6.5, followed by TENOVER FC A with an 8-year citation burst with 5.86 burst strength. OKEKE IN had the strongest burst 11.6 among cited authors.

Figure 5, depicts the cooperation network among Nigerian institutions/organizations on AMR research, where the nodes represent the institutions/organizations, and the sizes denote number of publications. The lines between two institutions/organizations denote that they

cooperate. The density of the links of an institutions/organization indicates the cooperation degree. The denser of the lines, the more collaboration for one institutions/organization with other institutions/organization. From the figure, University of Ibadan (n =366, links = 1586), University of Lagos (n =218, links = 1079), University of Nigeria (n =172, links = 430), Ahmadu Bello University (n =152, links = 647), Obafemi Awolowo University (n =141, links = 566) and Nigerian Institute of Medical Research (n =132, links = 883) have more lines. Therefore, they have more cooperation with other institutions/organizations.

3. AMR research themes and trends in Nigeria

Keywords are the condensation and reaction to the main content of the article, which can reflect the hot topic and the development trend related to the research field. We ran keyword co-occurrence network analysis with VOS viewer and arrived with a graph with 953 nodes and 6 Clusters (**Figure 6a**). The keywords “*Escherichia coli*” and “*Staphylococcus aureus*” displayed the highest frequency of 881 and 540, respectively. Followed by the keywords “ESBL” (470), “*Salmonella*” (468), “tetracycline” (465) and “plasmid” (412). Thematic research group of focus by Nigeria authors are; antimicrobial stewardship (25.50%), clinical and laboratory (25.39%), public health implication (23.92%), traditional and molecular methods (16.79%) and phytomedicine and drug discovery (8.39%). Additional insights on the contents of the clusters are described below;

a) Antimicrobial stewardship theme-based cluster (red, n= 243 items, 25.50%)

‘Antimicrobial stewardship’, ‘antibiotic prescription’, ‘antimicrobial resistance’, ‘aquatic environment’, ‘antimicrobial usage’, ‘attitude’, ‘awareness’, ‘burden’, ‘compliance’, ‘combination’, ‘evaluation’, ‘knowledge’, ‘middle income country’, ‘misuse’, ‘monitoring’, ‘mortality’, ‘observation’, ‘perception’, ‘point prevalence’, ‘practice’, ‘prescription’, ‘rational use’, ‘self-medication’, ‘standard’, ‘surveillance’, ‘survey’, ‘Systematic review’, ‘world health organization’ etc.

b) Clinical and laboratory theme-based cluster (green, n= 242 items, 25.39%)

‘Patients’, ‘outpatient’, ‘treatment’, ‘case’, ‘*Acinetobacter baumannii*’, ‘30ug’, ‘Adult’, ‘age group’, ‘*Aeromonas hydrophilia*’, ‘amoxicillin’, ‘amoxiclav’, ‘ampicillin’, ‘antibiotic’, ‘antenatal

clinic', 'antibiogram', 'antibiotics resistance profile', 'antibiotic susceptibility test', 'antibiotic resistance', 'Augmentin', 'azithromycin', 'bacillus', 'bacteriaemia', 'bacteria isolate', 'bacteria pathogen', 'bacterial infection', 'bacterial resistance', 'biofilm', 'blood agar', 'cefepime', 'cefixime', 'cephalosporin', 'cerebrospinal fluid', 'choice', 'cloxacillin', 'conventional biochemical test', 'cotrimoxazole', 'disk diffusion method', 'drug resistance', '*Enterobacter* spp', 'first line drugs', 'gram stain', 'high resistance', 'high vaginal swab', 'erythromycin', 'intensive care unit', '*Klebsiella*', 'Kirby Bauer disc diffusion method', 'levofloxacin', 'microbiology laboratory', 'multi drug resistant', 'pattern', 'otitis medium', '*Pseudomonas* spp', '*Shigella* spp', 'standard microbiological techniques', 'samples', 'UTI', 'wound infection', 'swab' etc.

c) Public health implication theme-based cluster (blue, n= 228 items, 23.92%)

'Isolate', 'strain', 'gene', 'animal', 'human', 'Abattoir environment', 'antibiotics resistance gene', 'animal production', 'antimicrobial residue', 'aquaculture', 'biofilm formation', 'cattle', 'coliform', 'environmental sample', 'faecal sample', 'ESBL', 'farm', 'fluoroquinolones', 'food handlers', 'gastroenteritis', 'indiscriminate use', 'livestock', '*Listeria spp*', 'meat sample', 'MLST', 'market', 'oxytetracycline', 'phenotypic characterization', 'poultry environment', 'pork', 'public health implication', 'raw meat', 'reservoir', 'resistance profile', '*Salmonella sp.*', 'salmonellosis', 'seafood', 'serotyping', 'whole genome sequencing', 'standard bacteriological method', '*Vibrio* spp', 'significance', 'virulence gene', 'resistance genes' etc.

d) Traditional and molecular methods cluster (yellow, n= 160 items, 16.79%)

'Molecular characterization', 'resistance gene', 'molecular detection', 'multiple antibiotics resistance gene', 'amp gene', 'Betalactamase', '*blaZ*', '*ctxm*', '*bla-shv*', '*bla-tem*', 'PCR', 'carbapenem', 'cefotaxime', 'CIP', 'clinical infection', 'clinical isolate', 'clinical sample', 'detection', 'plasmid profile', 'plasmid', '*pmqr*', 'sequencing', 'dissemination', 'double disc synergy test', '*E. coli*', 'ESBL', 'Gram negative', 'integrons', 'laboratory standard institute', 'MDR', 'MARI', '*mecA*', 'gene', 'pathogen of public health concern', 'methicillin resistant '*Staphylococcus aureus*', 'nasal swab', 'vancomycin', 'carriage', etc.

e) Phytomedicine and drug discovery themed cluster (purple, n= 80 items, 8.39%)

'Plant extract', 'aqueous extract', 'antibacterial activity', '*Candida spp*', 'concentration', 'efficacy', 'ethanol', 'flavonoid', 'Leafe', 'methanol', 'medicinal plants', 'microbial isolate', 'microorganisms', 'phytochemical', 'range', 'spectrum', 'tannin', 'discovery', 'technology', 'saponin', etc.

Figure 6b shows a Network map of the topics trend based on the most frequently used keywords. The current and future trends are highlighted in yellow.

Top cited articles with high field citation ratio (FCR) and relative citation ratio (RCR)

This study also revealed the top cited AMR articles indexed in Dimension database and sorted by FCR (**Table 2**). The FCR of the top 20 articles ranged from 4.09 to 36.54. The RCR ranged from 1.01 to 9.42. **Table S2** shows the Nigeria's top scholarly contributions to AMR indexed in Scopus database.

Table 1. Top 25 authors and cited authors in AMR research with the strongest citation bursts (sorted by beginning year).

Authors	Strength	Begin	End	1972 - 2022
OLUKOYA D	5.65	1988	1996	
COKER A	4.54	1992	2007	
BAKARE R	6.31	1995	2002	
OGUNSOLA F	4.45	2002	2016	
ADELOWO O	4.81	2008	2020	
OLOWE O	5.91	2012	2018	
ABIMIKU A	5.47	2012	2019	
UMOH VJ	4.88	2013	2015	
KWAGA J	4.63	2013	2022	
AKINDUTI P	4.21	2014	2020	
ANYANWU M	5.69	2015	2019	
OJO O	4.78	2015	2022	
ODUYEBO O	6.62	2016	2022	
ILIYASU G	4.23	2016	2022	
MOHAMMED Y	4.23	2016	2022	
EHINMIDU J	4.3	2017	2020	
IDOWU E	4.18	2017	2022	
CHAH K	4.63	2018	2022	
MUKHTAR M	4.02	2018	2022	
NGWAI Y	8.7	2019	2020	
OKEKE I	7.27	2019	2022	
IROHA I	4.8	2019	2022	
OLAYINKA B	8.53	2020	2022	
ODETOYIN B	4.26	2020	2022	
ADEBOWALE O	3.96	2020	2022	
Top Cited Authors				
Cited authors	Strength	Begin	End	1972 - 2022
AKO-NAI AK	6.5	1999	2017	
MENKIR A	6.92	2007	2012	
OKEKE IN	11.6	2009	2013	
TENOVER FC	5.86	2009	2017	
GHEBREMEDHIN B	7.0	2011	2014	
PATERSON DL	7.63	2012	2017	
LAWSON L	6.82	2013	2019	
SHITTU AO	5.86	2013	2018	
BAUER AW	5.29	2013	2014	
LIVERMORE DM	6.43	2014	2016	
AIBINU I	6.82	2015	2020	
ALTSCHUL SF	5.65	2015	2018	
MAGIORAKOS A-	9.44	2016	2022	
OJO OE	7.07	2017	2020	
ADESOKAN HK	5.75	2018	2022	
ADELOWO OO	5.63	2018	2022	
OLOSNO	7.93	2019	2022	
IBRAHIM SS	6.56	2019	2022	
ALHAJI NB	6.01	2019	2022	
KARIUKI S	5.98	2019	2020	
KRUMPERMAN PH	5.97	2019	2022	
IGBINOSA EO	5.86	2019	2022	
ABUBAKAR U	7.96	2020	2022	
AWORH MK	6.91	2020	2022	
ASHLEY EA	5.84	2020	2022	

Notes: The Blue bars mean the reference had been published; the red bars = Citation burstness (The start and end of the burst period is consistent with the length of red segment).

Table 2. Top 20 scholarly contributions to AMR authored by a Nigerian (Sorted by FCR).

Authors	DOI	Year	Times cited	RCR	FCR
Stewardson et al	10.1016/s1473-3099(18)30792-8	2019	92	8.45	36.54
Olaitan et al	10.1016/j.ijantimicag.2014.07.020	2014	203	9.42	24.66
Nurjadi et al	10.1093/jac/dku174	2014	75	3.13	19.42
Shittu et al	10.1186/1471-2180-11-92	2011	123	4.36	17.65
Ghebremedhin et al	10.1128/jcm.00648-09	2009	119	4.11	17.11
Kesah et al	10.1046/j.1469-0691.2003.00531.x	2003	96	2.22	16.92
Adesokan et al	10.4102/ojvr.v82i1.816	2015	61	3.37	14.57
Fashae et al	10.3855/jidc.909	2010	79	2.56	11.58
Aibinu et al	10.1111/j.1469-0691.2011.03730.x	2011	58	2.49	11.44
Marin et al	10.1371/journal.pntd.0002049	2013	82	3.23	9.96
Titilawo et al	10.1016/j.scitotenv.2015.03.095	2015	61	2.67	8.77
Ogbolu et al	10.1089/jmf.2006.1209	2007	77	1.01	8.09
Ogbolu et al	10.1016/j.ijantimicag.2010.08.019	2010	70	2.76	6.92
Titilawo et al	10.1007/s11356-014-3887-3	2015	63	2.73	6.21
Akinyemi et al	10.1016/j.puhe.2004.04.009	2005	53	1.15	5.98
Okeke et al	10.3201/eid0604.009913	2000	148	1.47	5.85
Fortini et al	10.1093/jac/dkr085	2011	71	2.76	5.69
Chigor et al	10.3390/ijerph7103831	2010	53	0.97	5.32
Iwalokun et al	10.1089/jmf.2004.7.327	2004	83	1.41	5.2
Soge et al	10.1093/jac/dki429	2005	64	1.83	4.09

Figure 1. Nigeria's annual contribution to AMR publication growth.




Figure 2. The top authors and journals of AMR research in Nigeria.





Figure 3. Centrality of top Universities (a) and Authors (b). Notes: each node in the network represents cited Author /University, the number of papers published by the author is represented by the size of the nodes, and the stroked rings represent the centrality index.



Figure 2: The co-authorship map of 1000 authors of authors in AMR research domain (T≥5). Notes: The size of node reflects the author’s co-occurrence frequencies, the link specify the co-occurrence association between authors, and the same colour of node represent the same cluster. In the figure there are 74 clusters ,7063 links and a total link strength of 10240.



researchers should participate actively in studies on AMR burden and contribute important scientific input to this field.

African Journal of Clinical and Experimental Microbiology was ranked as the top journal, publishing AMR documents with a minimum of one Nigerian institutional affiliation. The same journal was also the most cited in terms of published documents. Although Nigerian journals were among the top publishers of AMR research featuring at least a Nigerian author, there is a need to invest in strengthening local journals in Nigeria in order to achieve comparable indexing status and scientific standards to those of the world's top journals.

The frontier in a research area depicts the current growth path of a discipline in bibliometrics, and thus the references in the frontier article represent the field's intellectual core[28]. We described Nigeria's scholarly core in the field of AMR by mapping references. Themes on antimicrobial stewardship, clinical and laboratory practice, public health implication, traditional/molecular methods, and phytomedicine and drug discovery are major research trends of Nigeria's scientific community. These thematic distributions align with the observations of [29]. It is pertinent to boost research in all thematic areas of AMR research in the country to significantly facilitate processes that will slow down the spread of AMR in the country. Research themes exploring the use of molecular and artificial intelligence (AI) systems for AMR detection, surveillance, and drug discovery must be enhanced to confront rising bacterial resistance by significantly reducing the time taken to find new antimicrobials, improving diagnostic and treatment precision, as well as lowering costs [30].

A hotspot is a scientific topic of discussion in a series of documents which are inextricably connected to a particular time period[31]. The keywords are generalizations of the literature's topic[31]. A high-frequency keyword analysis can be used to identify hotspots in the field of AMR research. From our results "*Escherichia coli*" and "*Staphylococcus aureus*", "ESBL", "*Salmonella*", "tetracycline" and "Plasmid" were the most frequently used keywords. Nigerian researchers could conduct advanced studies based on the frequently discussed topics in this field.

An FCR range of 4.09 to 36.54 reflect the moderate to high scientific influence of Nigerian researchers in AMR research field. The theme of the top-cited documents in both the dimension and Scopus databases discussed are similar. The articles were focused on key AMR issues and determinants of human and animal importance relevant to the Nigerian and African population. Including; infectious diseases, Clinical and community health. Specifically, they highlighted the prevalence and/or Emergence of colistin resistance *Klebsiella pneumoniae*, Community-associated methicillin-resistant *Staphylococcus aureus*, carbapenem resistance, Enterobacteriaceae, multidrug resistant non-o1/non-o139 *Vibrio cholerae*, trimethoprim resistance gene *dfrG* in *Staphylococcus aureus*, plasmid-mediated quinolone resistance and betalactamases in *Escherichia coli*, CTX-M-15 extended-spectrum Beta-lactamase from *Klebsiella pneumoniae* etc. The identification of top-cited articles is useful as it reflects the diverse scope and health research fields that are of high interest to the readership of AMR, in addition to helping researchers identify the authors and institutions that have highly contributed to such influential studies for future research collaborations.

We are unlikely to witness a return to the heyday of antibiotic discovery, when numerous chemotypes were being regularly discovered and brought into clinical use in a short time frame. What does the antibiotic future hold? An assortment of approaches to infection treatment and prevention will have to be offered in the future. Anti-virulence strategies, small-molecule antibiotics, peptide-based antibiotics, bacteriophage monoclonal antibodies, probiotic strategies and vaccines are future trends in AMR and drug discovery.

Increasingly, the traditional Waksman platform returns recognized molecules. with recent breakthroughs in analytical chemistry, synthetic biology, genome mining, and genomics, the spectacular chemical diversity of nature can be taken advantage of by moving beyond the traditional Waksman platform. For instance, many environmental bacteria like actinomycetes harbour genetic programs to produce 20 to 40 natural products, while fungi can contain even more[32]. Avoiding actinomycetes bacteria is one strategy, for example. Some studies have identified 2 new antibiotics with new mechanisms of action, teixobactin, from the betaproteobacterium

Eleftheria terrae isolated from a soil specimen[33], and darobactin, from the gamma proteobacterium *Photorhabdus khaini* isolated from the gut of insect-killing nematodes[34].

The Myers laboratory has developed innovative modular pathways for the synthesis of new therapeutic tetracycline, macrolide, and lincosamide antibiotics, significantly expanding the diversity of these compound classes well beyond limits of natural biosynthesis[32]. Artificial intelligence approaches could also mine chemical libraries for therapeutic potential in order to develop new synthetic scaffolds[30].

Despite the difficulties in bringing peptide-based antibiotics to the clinic (most suffer from haemolytic activity or nephrotoxicity), optimism remains, and many AMPs are in various stages of development, targeted toward a range of pathogens[32].

Though naturally narrow in spectrum and must be combined with antibiotics, Antivirulence strategies aimed at abrogating pathogen virulence are a future trend. Sibofimloc, which prevents pathogenic *E. coli* from adhering to host epithelia is entering phase 2 trials[35].

Adjuvants or antibiotic combinations, such as b-lactamase inhibitors, can aid in the fight against resistance. A combo of bedaquiline, pretomanid, and linezolid, for instance, was recently approved to treat extensively drug - resistant *M. tuberculosis*, and more combinations are being studied[36].

In current history, the effectiveness and narrow spectrum of phage have been acknowledged as a valuable resource in the treatment of highly drug-resistant infections. The concept of using bacterial phages to treat drug-resistant infections is becoming more appealing. Phage therapy is increasingly being viewed as an alternative to conventional antibiotics in medicine, spurring more clinical trials, the majority of which are in phases one and two [32].

There are currently only 3 approved antimicrobial Monoclonal antibodies (mAbs), two of which inhibit the virulence of Bacillus anthracis (raxibacumab and obiltoxaximab), and one of which inhibits the virulence of *Clostridium difficile* (bezlotoxumab), reducing the incidence of reoccurrence. Only bezlotoxumab withstood clinical trials in its entirety[37], whereas the raxibacumab and obiltoxaximab were approved under the animal rule for biothreat organisms. But besides their

proven effectiveness, few are now in late-stage clinical trials (phase 2, *P. aeruginosa*: panobacumab/Aerumab; phase 2, *S. aureus*: omodenbamab/514G3 and; phase 3, *S. aureus*: suvrattoxumab and tosattoxumab[32].

Limitations

The data of this study were mainly derived from the dimensions database on August 20th 2022, and other literature databases like Web of Science, Open Alex and Semantic scholar were not searched. Therefore, we might not have been able to retrieve all papers on this topic, and selection bias may have been present.

Conclusion and recommendations

According to this research, Nigerian institutions and organizations contributed meagerly (0.2%) to AMR research output globally. There is a pressing need to strengthen the country's research capacity and expedite the use of insights provided by its scientists across all subject areas in order to make a significant contribution to AMR response. Although this response is not always tied to local data but instead relies on adapting global policies and those of global health organizations, developing local data to better understand AMR trends is still an excellent method for Nigeria to respond to health emergencies. Nigeria's potential to respond to health issues needs to be strengthened, and investing in research is one excellent way to do so.

Availability of data and materials

The raw datasets generated are accessible upon demand to the corresponding authors (suleykestler2@gmail.com, shuaibusuleiman60@yahoo.com)

Competing interest

We declare that we have no conflict of interest.

Funding

No funding was received.

References

- 1-**Prestinaci F, Pezzotti P, Pantosti A.** Antimicrobial resistance: a global multifaceted phenomenon. *Pathog Glob Health* 2015; 109: 309–318.
- 2-**World Health Organization (WHO).** Antimicrobial resistance: global report on surveillance. World Health Organization, 2014.

- 3-**Chukwu EE, Oladele DA, Awoderu OB, et al.** A national survey of public awareness of antimicrobial resistance in Nigeria. *Antimicrob Resist Infect Control* 2020; 9: 72.
- 4-**Achi CR, Ayobami O, Mark G, Egwuenu A, Ogbolu D, Kabir J.** Operationalising One Health in Nigeria: Reflections From a High-Level Expert Panel Discussion Commemorating the 2020 World Antibiotics Awareness Week. *Front Public Health* 2021; 9: 673504.
- 5-**Harant A.** Assessing transparency and accountability of national action plans on antimicrobial resistance in 15 African countries. *Antimicrobial Resistance & Infection Control* 2022; 11: 15.
- 6-**Knobler SL, Infections I of M (US) F on E, Lemon SM, et al.** Factors Contributing to the Emergence of Resistance. National Academies Press (US), <https://www.ncbi.nlm.nih.gov/books/NBK97126/> (2003, accessed 17 September 2022).
- 7-**World Health Organization (WHO).** Experts caution Nigerians against indiscriminate use of antibiotics to reduce future health crisis. World Health Organization / *Regional Office for Africa*, <https://www.afro.who.int/news/experts-caution-nigerians-against-indiscriminate-use-antibiotics-reduce-future-health-crisis> (accessed 17 September 2022).
- 8-**Auta A, Banwat SB, David S, Dangiwa DA, Ogbolu E, Tor-anyiin AJ.** Antibiotic Use in Some Nigerian Communities: Knowledge and Attitudes of Consumers. *Tropical Journal of Pharmaceutical Research* 2013; 12: 1087–1092.
- 9-**Stivers T.** Managing Patient Pressure to Prescribe Antibiotics in the Clinic. *Paediatr Drugs* 2021; 23: 437–443.
- 10-**Adesokan HK, Akanbi IO, Akanbi IM, Obaweda RA.** Pattern of antimicrobial usage in livestock animals in south-western Nigeria: The need for alternative plans. *Onderstepoort J Vet Res* 2015; 82: 816.
- 11-**Kotwani A, Joshi J, Kaloni D.** Pharmaceutical effluent: a critical link in the interconnected ecosystem promoting antimicrobial resistance. *Environ Sci Pollut Res* 2021; 28: 32111–32124.
- 12-**Sinha R, Shukla P.** Antimicrobial Peptides: Recent Insights on Biotechnological Interventions and Future Perspectives. *Protein and Peptide Letters* 2019; 26: 79–87.
- 13-**Liang C, Luo A, Zhong Z.** Knowledge mapping of medication literacy study: A visualized analysis using CiteSpace. *SAGE Open Medicine* 2018; 6: 205031211880019.
- 14-**Ding X, Yang Z.** Knowledge mapping of platform research: a visual analysis using VOSviewer and CiteSpace. *Electron Commer Res* 2022; 22: 787–809.
- 15-**Adeiza S, Shuaibu A.** Trends in monkeypox research: A sixty-year bibliometric analysis. *Microbes and Infectious Diseases* 2022; 3: 500–513.
- 16-**Li Y, He M, Zou Z, Bian X, Huang X, Yang C, et al.** Artificial liver research output and citations from 2004 to 2017: a bibliometric analysis. *PeerJ* 2019; 6: e6178.
- 17-**Hook DW, Porter SJ, Herzog C.** Dimensions: Building Context for Search and Evaluation. *Front Res Metr Anal* 2018; 3: 23.
- 18-**Van Eck N, Waltman L.** Software survey: VOSviewer, a computer program for bibliometric mapping. *scientometrics* 2010; 84: 523–538.
- 19-**Perianes-Rodriguez A, Waltman L, van Eck NJ.** Constructing bibliometric networks: A comparison between full and fractional counting. *Journal of Informetrics* 2016; 10: 1178–1195.

- 20-**Huang T, Wu H, Yang S, Su B, Tang K, Quan Z, et al.** Global Trends of Researches on Sacral Fracture Surgery: A Bibliometric Study Based on VOSviewer. *Spine* 2020; 45: E721.
- 21-**Suleiman AS, Abdulmalik SB, Ghazali SM.** Extended Spectrum Betalactamase Research mapping; a 32-year Temporal and geographical perspective. *Epub ahead of print* 2 August 2022. DOI: 10.21203/rs.3.rs-1898374/v1.
- 22-**Chen C.** CiteSpace II: Detecting and visualizing emerging trends and transient patterns in scientific literature. *J Am Soc Inf Sci* 2006; 57: 359–377.
- 23-Dimensions, **what is the FCR?** How is it calculated? <https://dimensions.freshdesk.com/support/solutions/articles/23000018848-what-is-the-fcr-how-is-it-calculated-> (accessed 17 September 2022).
- 24-**Sweileh WM, Moh'd Mansour A.** Bibliometric analysis of global research output on antimicrobial resistance in the environment (2000–2019). *Global Health Research and Policy* 2020; 5: 37.
- 25-**Torres RT, Carvalho J, Cunha MV, Serrano E, Palmeira JD, Fonseca C.** Temporal and geographical research trends of antimicrobial resistance in wildlife—a bibliometric analysis. *One Health* 2020; 11: 100198.
- 26-**Payumo J, Alocilja E, Boodoo C, Luchini-Colbry K, Ruegg P, McLamore E, et al.** Next Generation of AMR Network. *Encyclopedia* 2021; 1: 871–892.
- 27-**Leydesdorff L.** Betweenness centrality as an indicator of the interdisciplinarity of scientific journals. *Journal of the American Society for Information Science and Technology* 2007; 58: 1303–1319.
- 28-**Qi S, Hua F, Xu S, Zhou Z, Liu F.** Trends of global health literacy research (1995–2020): Analysis of mapping knowledge domains based on citation data mining. *Plos one* 2021; 16: e0254988.
- 29-**Luz CF, van Niekerk JM, Keizer J, Beerlage-de Jong N, Braakman-Jansen LMA, Stein A, et al.** Mapping twenty years of antimicrobial resistance research trends. *Artificial Intelligence in Medicine* 2022; 123: 102216.
- 30-**Rabaan AA, Alhumaid S, Mutair AA, Garout M, Abulhamayel Y, Halwani MA, et al.** Application of Artificial Intelligence in Combating High Antimicrobial Resistance Rates. *Antibiotics* 2022; 11: 784.
- 31-**Wang Z, Ma D, Pang R, Xie F, Zhang J, Sun D.** Research Progress and Development Trend of Social Media Big Data (SMBD): Knowledge Mapping Analysis Based on CiteSpace. *IJGI* 2020; 9: 632.
- 32-**Cook MA, Wright GD.** The past, present, and future of antibiotics. *Sci Transl Med* 2022; 14: eabo7793.
- 33-**Ling LL, Schneider T, Peoples AJ, poering AL, Engels I, Conlon BP, et al.** A new antibiotic kills pathogens without detectable resistance. *Nature* 2015; 517: 455–459.
- 34-**Imai Y, Meyer KJ, Iinishi A, Favre-Godal Q, Green R, Manuse S, et al.** A new antibiotic selectively kills Gram-negative pathogens. *Nature* 2019; 576: 459–464.
- 35-**Reinisch W, Hébuterne X, Buisson A, Schreiber S, Desreumaux P, Primas C. et al.** Safety, pharmacokinetic, and pharmacodynamic study of sibofimloc, a novel FimH blocker in patients with active Crohn's disease. *Journal of Gastroenterology and Hepatology.*

36-**Wright GD**. Antibiotic adjuvants: rescuing antibiotics from resistance. *Trends in microbiology* 2016; 24: 862–871.

37-**Allio T**. The FDA Animal Rule and its role in protecting human safety. *Expert Opinion on Drug Safety* 2018; 17: 971–973.

Adeiza SS, Shuaibu MG, Shuaibu AB. Knowledge mapping of Nigeria's scientific contribution to antimicrobial resistance research: A visualized investigation using VOS viewer and cite space. *Microbes Infect Dis* 2024; 5(2): 588-602.