

Original article

Antibiotic resistance pattern of *Staphylococcus aureus* isolated from nostrils of healthy undergraduates of Madonna University Elele Campus, Rivers State, Nigeria

Franklin Ifechukwu Onyeka ¹, David Chinemerem Nwobodo ^{2,3*}, Ignatus Chinedu Umenne ⁴, Emmanuel Edet Atada ¹, Chibundu Amalachukwu Ojukwu ¹, Maureen Adaobi Aniekwe ¹, Jeffrey James Philomena ¹, Joseph Chinedu Ikem ^{1,3}

1- Department of Pharmaceutical Microbiology & Biotechnology, Madonna University, Elele, River State.

2- Department of Microbiology, Renaissance University, Enugu.

3- Department of Pharmaceutical Microbiology & Biotechnology, Nnamdi Azikiwe, University, Nigeria.

4- Department of Microbiology, Micheal Okpara University of Agriculture, Umudike, Abia State.

ARTICLE INFO

Article history:

Received 16 May 2020

Received in revised form 25 June 2020

Accepted 26 June 2020

Keywords:

Staphylococcus aureus

Nasal

Antibiotics

Resistance

Nigeria

ABSTRACT

Background: Antimicrobial resistance is a global health issue with particular concern in developing countries, where there are poor antibiotic regulatory policies. *Staphylococcus aureus* (*S. aureus*) has become a persistent nosocomial and community-acquired pathogen that has become a global menace. This study determined the prevalence and resistance patterns of nasal *S. aureus* isolated from healthy students of Madonna University, Elele Campus, Nigeria. **Methods:** One hundred and fifty nasal specimens of healthy students were cultured and screened for *S. aureus* using standard microbiological protocols and their antibiotic profile susceptibility was investigated using the disc diffusion method according to the Clinical and Laboratory Standards Institute on Mueller Hinton agar. **Results:** A total of 78 (52%) *S. aureus* isolates were obtained from 150 specimens screened. The pattern of *S. aureus* resistance to the antibiotics varied in prevalence by agent in the descending order as follows; cefuroxime = ceftazidime > cloxacillin > erythromycin > augmentin > ofloxacin > ceftriaxone > gentamicin. The isolates showed an overall 100% resistance to ceftazidime and cefuroxime. Gentamicin, ceftriaxone, and ofloxacin recorded a susceptibility rate of 98.7%, 89.7%, and 79.5% respectively. **Conclusion:** This study reveals a high prevalence of *S. aureus* colonization in students in the study area. Also, these isolates could still be treated with gentamicin, ceftriaxone, and fluoroquinolones. However, there is increasing resistance to commonly prescribed antibiotics. There is a need for stringent strategies in the control of antibiotic misuse and resistance.

Introduction

Staphylococcus aureus (*S. aureus*) is a Gram-positive, coagulase-positive bacterium found on the skin and in the nose of most healthy people [1]. The nose is one of the few openings that bacteria have free passage to get inside the body. Thus, the nose and nasal passages can be a perfect environment for some bacterial communities to thrive. *Staphylococcus*

aureus is known as normal flora in the skin and occasionally carried by over 60% of the world's population during their lifetime, without it causing any harm [2]. However, it is considered an opportunistic pathogen for humans and animals if found in the bloodstream and tissues. Over the past decades, *S. aureus* has been recognized as one of the most

important bacterial pathogens which significantly contribute to hospital and community-acquired infections all over the world [1,3-5]. It is found to cause various illness ranging from minor skin infections such as pimples, impetigo, boils, cellulitis, scalded skin syndrome, folliculitis, furuncles, carbuncles, and abscesses, to life-threatening diseases such as pneumonia, osteomyelitis, meningitis, Toxic Shock Syndrome, endocarditis and septicemia [4].

Infections caused by *S. aureus* are worrisome, because of its intrinsic virulence, its ability to adapt to various environmental conditions, and ease in the acquisition of resistance to new drugs [6]. Recently, many isolates of *S. aureus* have evolved, and displayed resistance to traditional antimicrobial chemotherapy and their prevalence outside the hospital is of potential epidemiological threat. Resistance to commonly available and affordable antibiotics pose a major concern in the management of bacterial infections, especially in developing countries [7,8]. Lack of concern or good judgment in the use of antibiotics in human medicine, and for prophylaxis in animal husbandry contribute greatly to the emergence of multidrug-resistant (MDR) strains.

In several studies worldwide, *S. aureus* from normal flora seems to constitute an important reservoir of antimicrobial resistance genes [1,8-10], which can be transferred to other microbial pathogens, thus, propagating the resistance traits among microbial populations [11]. The prevalence of antibiotic-resistant *staphylococci* at various skin sites in both healthy and hospitalized patients have received much attention because of the role of these organisms as nosocomial pathogens, especially in an immunocompromised host [8].

The World Health Organization (WHO) in 2014, reported significant gaps in antibiotics surveillance, absence of standards methodology, data sharing, and coordination. According to the report, major gaps were identified in Africa, South-East Asia, and the Eastern Mediterranean regions [12]. Data on the frequency of nasal carriage, and *S. aureus* resistance to antibiotics in Elele is not known. The indiscriminate use of antibiotics before consulting a physician for a prescription is common in Nigeria. Students are major victims, as they prefer to self-prescribe or buy cheap over-the-counter drugs, and not see a physician due to financial constraints. On the other hand, most physicians prescribe and treat patients with broad-spectrum antibiotics before or without microbiological investigations [13]. These factors are key players in the development and spread

of bacterial resistance to affordable and commonly used antibiotics. Antibiotic resistance surveillance data is necessary to better inform the effective administration of antibiotic therapy, and further reduce the spread of drug resistance. Thus, this study was designed to determine the prevalence and antibacterial susceptibility profile of *S. aureus* isolated from nostrils of apparently healthy undergraduates in Elele, South-Southern Nigeria. Our results may improve antibiotic-use decisions/policies.

Materials and Methods

Sample collection

This study was conducted in the Department of Pharmaceutical Microbiology and Biotechnology Laboratory, Madonna University Elele Campus, River State. A total of one hundred and fifty (150) nasal samples were randomly obtained with sterile swab sticks from healthy undergraduates students of Madonna University Elele Campus (**Figure 1**), Rivers State, Nigeria. An informed consent was obtained from each participant. The population comprised undergraduates, all aged between 18 and 28 years. All sampling procedures were according to the guidelines of the National Health Research Ethics Committee, Nigeria (www.nhrec.net).

Isolation and identification of *S. aureus*

Swab samples were cultured on Mannitol salt agar (Oxoid, England) and incubated for 18-24 hours at 37 °C. The pure culture of each isolate was obtained by the streak plate technique. Identification of isolates was performed by colony morphology, Gram staining, catalase test, and coagulase test [1]. Gram-positive, catalase, and coagulase-positive isolates were considered as *S. aureus* [1,14].

Antibiotic susceptibility testing (AST)

Antibiotic susceptibility of the isolates was determined using the modified Kirby-Bauer disc diffusion technique [11]. The isolates were briefly cultured in Mueller Hinton broth at 37 °C for 24 hours. A suspension of each isolate standardized to match 0.5 McFarland were inoculated onto sterile agar plates. The plates were allowed to set and the antibiotic sensitivity disc (ABTEK, India) containing; ceftazidime (CAZ) 30µg, cefuroxime (CRX) 30µg, gentamicin (GEN) 10µg, ceftriaxone (CTR) 30µg, erythromycin (ERY) 5µg, cloxacillin (CXC) 5µg, ofloxacin (OFL) 5µg, augmentin (AUG) 30µg. The plates were incubated at 37 °C for 24 hours and the resultant inhibition zone diameters (IZDs) were measured and recorded. The obtained results were

interpreted based on the guidelines of the Clinical Laboratory Standard Institute [15].

Statistical analysis

Data obtained were analyzed using Microsoft Excel (2016). Results were presented as descriptive statistics in terms of relative frequency and percentage.

Figure 1. Map of the geographical location of Elele Nigeria, the study area (Map data @2019 Google).



Results


The results show that seventy-eight (78) *S. aureus* strains were isolated from the one hundred and fifty (150) nasal samples screened, giving a prevalence rate of 52%. The antibiotic susceptibility pattern of *S. aureus* presented in **table (1)** shows that gentamicin, ceftriaxone, and ofloxacin had the best anti-staphylococcal activities, with only one (1) isolate resistant to gentamicin. 93% of the isolates were resistant to cloxacillin, while all the studied isolates

were completely resistant to ceftazidime and cefuroxime. The prevalence rate of *S. aureus* susceptibility pattern to antibiotics obtained in this study can be summarized in descending order as follows; gentamicin (98.7%) > ceftriaxone (89%) > ofloxacin (79.5%) > augmentin (64.1%) > Erythromycin (20.5%) > cloxacillin (6.4%) > cefuroxime (0%) = ceftazidime (0%). **Figure 2** represents a comparison of the susceptibility and resistance of the isolated *S. aureus* against various antibiotics used in the study.

Table 1. Antibiotic susceptibility pattern of *S. aureus* isolated from pharmacy students of Madonna University.

Antibiotics	Number susceptible	%	Number resistant	%
Gentamicin	77	98.7	1	1.3
Ceftriaxone	70	89.7	8	10.3
Ofloxacin	62	79.5	16	20.5
Augmentin	50	64.1	28	35.9
Erythromycin	16	20.5	62	79.5
Cloxacillin	5	6.4	73	93.6
Ceftazidime	0	0	78	100
Cefuroxime	0	0	78	100

Figure 2. Plates A and B showing the susceptibility patterns of isolated *S. aureus* to the selected antibiotics. *Staphylococcus aureus* cultured on both plates were remarkably susceptible to gentamicin, ceftriaxone, and augmentin.



Discussion

The nose is known to harbor numerous and diverse groups of microorganisms. *Staphylococcus aureus* is a common bacterium carried by most people in the nose and skin and is regarded as a major risk factor for the development of infections in people. Infections caused by *S. aureus* are commonly endogenous, meaning they are caused by strains already colonizing the patient [9]. *Staphylococcus aureus* strains have been reported to cause infections in healthy people, and these infections are increasing in various regions and countries [1,5].

The carrier rate of *S. aureus* in this study was 52% which is similar to that obtained in other studies carried out in Eastern Nigeria 50%, 53%, and 56.3% [8,16,17]. However, **Nsofor et al.** [18] reported a higher *S. aureus* prevalence rate (62.9%) among school pupils in the same locality (Elele) as the study area. These results suggest a high nasal carriage of *S. aureus* among people living and residing in Elele. The *S. aureus* in this study was found to be resistant to cefuroxime, cloxacillin, and ceftazidime, but sensitive to gentamicin, ceftriaxone, and ofloxacin. Gentamicin and ceftriaxone come as injectable while ofloxacin belongs to the fluoroquinolones family. Beta lactams including penicillin derivatives and cephalosporins are commonly used antibiotics freely available in open market and other unregistered drug outlets.

This observed resistance pattern is reflective of the rate of antibiotics misuse and abuse arising from self-medications, which is often associated with inadequate dosage and failure to comply with treatments. Also, the availability of antibiotics to consumers across the counters with or without prescription. Several other researchers have also reported *S. aureus* resistance to commonly used antibiotics in community settings [16,17,19].

The observed high sensitivity of *S. aureus* to fluoroquinolones (ofloxacin) and gentamicin in this study were also observed when antibiotic susceptibility tests were conducted on healthy school pupils in Agulu, South-eastern Nigeria [19]. **Nwankwo and Nasiru** [20] reported similar antibiotic susceptibility patterns of *S. aureus* from clinical isolates in a tertiary health institution in Kano, North-west Nigeria.

It is important to note that from the *S. aureus* resistance profile results, antibiotics showing relatively good susceptibility profile (gentamicin, ceftriaxone, and ofloxacin) are agents whose antimicrobial mode of actions is based on the inhibition of protein synthesis, inhibition of cell wall synthesis and inhibition of DNA Gyrase, and are known to be generally active against Staphylococcal infections [1,21].

The increasing frequency of antimicrobial resistance among undergraduates and people in Elele is of great concern to both healthcare providers and the public. This is because the high incidence of MDR *S. aureus* in a community will make treatment of these infections more difficult [10,22].

Conclusion

Our results show the urgent need to discourage antibiotics abuse in order to prevent MDR in community-acquired staphylococcal infections. On the other hand, health sectors should base their prescriptions on proper antibiotic susceptibility results. This is important to preserve the efficacy of antibiotic drugs, and thus contain the emergence and widespread of drug resistance.

Contributors

All authors participated equally in the design, research and preparation of this manuscript.

Authorship

All authors have approved the submission of this version of the manuscript and takes full responsibility for it.

Competing interest: None declared by authors.

Funding: There was no financial aid received.

References

- 1-**Anie CO, Ugwu MC, Ibezim EC, Esimone CO.** Antibigram of methicillin-resistant *Staphylococcus aureus* isolates among healthy human subjects in Oleh, South-Southern Nigeria. International Journal of Current Microbiology and Applied Sciences 2017; 6(9): 3710-3716.
- 2-**Raja Ram G, Prashanna M, Ganga GC.** Antibiotic resistance pattern of *Staphylococcus aureus* with reference to MRSA isolates from pediatric patients. Future Sci OA 2020; 6(4): FSO464.
- 3-**Yassin NA, Mohammed HH, Ahmad AM.** Antibigraming profiles of *Staphylococcus aureus* isolated from various clinical specimens in Duhok city Iraq. Advance Tropical Medicine and Public Health International 2013; 3(1): 25-31.
- 4-**Nsofor CA, Nwokenkwo VN, Ohale CU.** Prevalence and antibiotic susceptibility pattern of *Staphylococcus aureus* isolated from various clinical specimens in south east Nigeria. MOJ Cell Science and Report 2016; 3(2): 60-63.
- 5-**Ugwu MC, Anie CO, Ibezim EC, Esimone CO.** Antimicrobial evaluation of methicillin-resistant *Staphylococcus aureus* nasal carriage amongst healthy students in Agbor, Delta State, Nigeria. Archives of Clinical Microbiology 2016; 7(2):13.
- 6-**Karchmer AW.** From theory to practice: resistance in *Staphylococcus aureus* and new treatments. Clinical Microbiology and Infections 2006; 12(8): 15-21.
- 7-**Kaplan SL, Hulten KG, Gonzalez BE, Hammerman WA, Lamberth L, Versalovic J, et al.** Three-year surveillance of community-acquired *Staphylococcus aureus* infections in children. Clin Infect Dis 2005; 40(12):1785-1791.
- 8-**Ugwu MC, Ibezim EC, Odimegwu D.** Antibiotic resistance patterns of *Staphylococcus aureus* isolated from nostrils of healthy human subjects in a South-Eastern Nigeria Locality. Macedonian Journal of Medical Sciences 2009; 2(4):294-300.
- 9-**Al-Haj NA, Hauter JM, Al-Bulili NH, Al-Hotami RA, Al-Horaibi MT.** Nasal carriage of among students of public schools in Sana'a, Yemen. Research Journal of Microbiology 2018; 13(1): 65-69.
- 10-**Tekalign K, Bacha K.** Prevalence and antibiotic susceptibility pattern of methicillin-resistant *Staphylococcus aureus* (MRSA) among primary school children and prisoners in Jimma Town, Southwest Ethiopia. Annals of Clinical Microbiology and Antimicrobials 2013; 12(11): 1-12.
- 11-**Van den Bogaard AE, Stobberingh EE.** Epidemiology of resistance to antibiotics. Links between animals humans. Int J Antimicrob Agents 2000; 14(4): 327-35.
- 12-**World Health Organization (WHO).** Antimicrobial resistance: global report on surveillance. World Health Organization 2014.
- 13-**Aboderin OA, Abdu AR, Odetoyin BW, Lamikanra A.** Antimicrobial resistance in *Escherichia coli* strains from urinary tract infections. Journal of National Medical Association 2009; 101(12): 1268-1273.
- 14-**Cheesbrough M.** Biochemical tests to identify bacteria. District Laboratory practices in tropical Countries part 2 2006; 62-70.

- 15-**Clinical Laboratory Standard Institute (CLSI)**. Performance standards for antimicrobial susceptibility testing; document M100 26th.
- 16-**Chigu CO, Ezeronye OU**. Antibiotic resistant *Staphylococcus aureus* in Abia State of Nigeria. African Journal of Biotechnology 2003; 2(10): 374-378.
- 17-**Nsofor CA, Nwokenkwo VN, Nwaokpa C**. Nasal carriage of *Staphylococcus aureus* among apparently healthy school children in Owerri, Nigeria. International Journal of Microbiology and Immunology Research 2015; 1(2): 16-21.
- 18-**Nsofor CA, Ezeh OI, Nwabia BC**. Nasal carriage of multi-drug resistant *Staphylococcus aureus* among apparently healthy school children in Elele, Nigeria. MOJ Cell Science and Report 2013; 1(2): 117-120.
- 19-**Ugwu MC, Mokwe N, Ejikeugwu PC, Gugu TH, Enemor EC, Eze CO, et al**. Antibigram of *Staphylococcus aureus* from healthy school pupils in Agulu, Southeastern Nigeria. Int Jou of Res in Pharma and Bioscs 2015; 2(4): 5-9.
- 20-**Nwankwo EO, Nasiru M**. Antibiotic susceptibility pattern of *Staphylococcus aureus* from clinical isolates in tertiary health institution in Kano, North-Western Nigeria. Pan African Medical Journal 2011; 1937-8688.
- 21-**Leiboivitz E**. The use of fluoroquinolones in children. Curr Opin 2006; 18(1): 64-70.
- 22-**Rijal KR, Shrestha N, Pahari N**. Methicillin resistant *Staphylococcus aureus* in patients visiting western regional hospital, Pokhara. JIOM 2008; 30(1): 21–25.

Onyeka FI, Nwobodo D, Umenne IC, Atada EE, Ojukwu CA, Aniekwe MA, Philomena JJ, Ikem JC. Antibiotic resistance pattern of *Staphylococcus aureus* isolated from nostrils of healthy undergraduates of Madonna University Elele Campus, Rivers State, Nigeria. Microb Infect Dis 2021; 2 (2): 280-285.